

Interacting holographic dark energy

Winfried Zimdahl^{*1} and Diego Pavón^{†2}

¹ *Departamento de Física, Universidade Federal do Espírito Santo,
CEP29060-900 Vitória, Espírito Santo, Brazil*

² *Departamento de Física, Facultad de Ciencias,
Universidad Autónoma de Barcelona,
08193 Bellaterra (Barcelona), Spain*

(Dated: February 5, 2008)

Abstract

We demonstrate that a transition from decelerated to accelerated cosmic expansion arises as a pure interaction phenomenon if pressureless dark matter is coupled to holographic dark energy whose infrared cutoff scale is set by the Hubble length. In a spatially flat universe the ratio of the energy densities of both components remains constant through this transition, while it is subject to slow variations for non-zero spatial curvature. The coincidence problem is dynamized and reformulated in terms of the interaction rate. An early matter era is recovered since for negligible interaction at high redshifts the dark energy itself behaves as matter. A simple model for this dynamics is shown to fit the SN Ia data. The constant background energy density ratio simplifies the perturbation analysis which is characterized by non-adiabatic features.

^{*} E-mail: zimdahl@thp.uni-koeln.de

[†] E-mail address: diego.pavon@uab.es

I. INTRODUCTION

Nowadays an overwhelming, direct and indirect, observational evidence supports the idea that the Universe is currently undergoing a phase of accelerated expansion. The most recent and precise confirmation of this rather unexpected feature (see, however [1, 2]) was provided by the 3rd year data of the WMAP mission [3]. Likewise, data from high-redshift supernovae type Ia [4], the cosmic microwave background radiation [5], the large scale structure [6], the integrated Sachs–Wolfe effect [7], and weak lensing [8], endorse it to the point that the discussion has now shifted to when the acceleration began and which might be the agent behind it. For recent reviews see [9].

According to our understanding on the basis of Einstein’s gravity, more than 70% of the cosmic substratum, dubbed dark energy, must be endowed with a high negative pressure. Less than 30% is found to be pressureless matter. Most of the latter is in the form of cold dark matter; only about 5% corresponds to “normal” baryonic matter. Currently, we neither know what the dark matter is made of nor do we understand the nature of dark energy.

Lacking a fundamental theory, most investigations in the field are phenomenological and rely on the assumption that these two unknown substances evolve independently, i.e., their energies are assumed to obey separate conservation laws. In particular, this implies that the dark matter energy density varies as a^{-3} , where a is the scale factor of the Robertson–Walker metric. The behavior of the density of the dark energy is then entirely governed by its equation of state, the determination of which is a major subject of current observational cosmology. One should realize, however, that any coupling in the dark sector will change this situation. A coupling will modify the evolution history of the Universe. As one of the consequences, the energy density of the (interacting) dark matter will no longer evolve as a^{-3} . Ignoring a potentially existing interaction between dark matter and dark energy from the start, may result in a misled interpretation of the data concerning the dark energy equation of state. Das *et al.* [10] and Amendola *et al.* [11] have shown that a measured phantom equation of state may be mimicked by an interaction, while the bare equation of state may well be of a non-phantom type. Further, models showing interaction fare well when contrasted with data from the cosmic microwave background [12] and matter distribution at large scales [13]. It can therefore be argued that the possibility of dark energy to be in interaction with dark matter must be taken seriously.

On the other hand, there exist limits for the strength of this interaction for various configurations [14]. It is characteristic for these approaches that they admit a non-interacting limit. If the interaction is switched off, they continue to represent models of a mixture of dark matter and dark energy. The best known examples are models with a decaying cosmological “constant”: if the decay is switched off, they reduce to the Λ CDM model. An interaction here leads to (possibly important) corrections of the non-interacting configuration. E.g., for a given (not necessarily constant) negative equation of state parameter of the dark energy, the interaction manifests itself in third order in the redshift in the luminosity distance of supernovae type Ia [15]. It does not, however, (directly) influence the leading orders.

The qualitatively new aspect of the present approach is the circumstance that an interaction is crucial already in leading order. In this context the accelerated expansion itself is a consequence of the interaction. The coupling does not just lead to higher order corrections of a non-interacting reference model which by itself provides an accelerated expansion of the Universe. The non-interacting limit of the present approach reproduces an Einstein-de Sitter universe, i.e., there is no accelerated expansion if the interaction is switched off. This non-interacting limit is supposed to characterize our Universe at high redshifts.

The data also suggest that the present Universe is nearly spatially flat [3], [16]. Many studies neglect the spatial curvature term and focus solely on the spatially flat case, thereby taking for granted that the spatial curvature term necessarily leads to trifling corrections. Several of the not so many works that retain that term, for the sake of generality, even seem to justify that it is of minor importance. On the other hand, thanks to the increasing observational precision also higher order corrections may become within reach in the not too distant future. The authors of [17] have demonstrated that the spatial curvature enters the luminosity distance of SNIa supernovae in third order in redshift (see also [18]). It is also known that there are degeneracies between the curvature and the dark energy equation of state in the corresponding parameter space [19]. The possible relevance of spatial curvature for the lowest multipoles in the cosmic microwave background radiation was discussed in [20]. These examples indicate that apart from general theoretical grounds, the observational situation will require the inclusion of spatial curvature, even if its contribution is small, at some level of precision. Further, as demonstrated by Ichikawa *et al.* [21], depending on the parametrization of the equation of state parameter the present value of the spatial curvature can be as large as 0.2. Therefore, the additional dynamics provided by the curvature, should

certainly not be dismissed too quickly since it contains information which is needed to further restrict the cosmological parameter space. In this connection, Clarkson *et al.* [22] have shown that by excluding the spatial curvature when interpreting the empirical data one can incur in gross mistakes when reconstructing the equation of state of dark energy. We mention that an effective spatial curvature term also appears as the result of an averaging procedure within the “macroscopic gravity” approach [23] (for a further discussion of the effects of averaging on cosmological observations see, e.g., [24]).

In this paper we consider pressureless dark matter in interaction with an unknown component which is supposed to describe dark energy. We neither specify the dark energy equation of state nor the interaction rate from the beginning. These two (generally time dependent) parameters will influence the ratio of the energy densities of dark matter and dark energy. The behavior of this ratio is crucial for the “conventional” form of the “coincidence problem”, namely: “why are the matter and dark energy densities of precisely the same order today”. In principle, matter and dark energy redshift at different rates. We show, that there exists a preferred class of dark energy models for which the dynamics of the energy density ratio is entirely determined by the spatial curvature. For vanishing curvature the energy density ratio remains constant. These models are singled out by a dependence $\rho_X \propto H^2$, where ρ_X is the dark energy density and $H = \dot{a}/a$ is the Hubble parameter. Exactly this dependence is characteristic for a certain type of dark energy models, inspired by the holographic principle [25, 26].

Holographic dark energy models must specify an infrared cutoff length scale [27]. The choice of this scale is presently a matter of debate. The most obvious choice, the Hubble length, seemed to be incompatible with an accelerated expansion of the Universe [28] (see, however, [29, 30]). This is why, starting with the work of Li [31], many researchers have adopted the future event horizon as the cutoff scale as this choice allows for a sufficiently negative equation of state parameter and hence an accelerated expansion -see, e.g. [32].

In a previous paper [33], we showed that a cutoff set by the Hubble length may well be compatible with an accelerated expansion provided that the dark energy and dark matter do not evolve separately but interact, also non-gravitationally, with each other. In this setting, a negative equation of state parameter that gives rise to accelerated expansion arises as a direct consequence of the interaction. Here we put this feature in a broader context and demonstrate that a constant or slowly varying (as the consequence of a non-vanishing

spatial curvature) energy density ratio is compatible with a transition from decelerated to accelerated expansion under the condition of a growing interaction parameter. We show that in holographic dark energy models with a Hubble length cutoff a transition from decelerated to accelerated expansion is realized as a pure interaction effect. At high redshifts and for negligible interaction the dark energy equation of state approaches the equation of state for matter, such that a preceding matter era is naturally recovered. In this context the coincidence problem can be rephrased as follows: “why is the interaction rate between dark matter and dark energy of the order of the Hubble rate precisely at the present epoch?” This reformulation allows us to address the coincidence problem as part of the interaction dynamics in the dark sector.

Different interaction rates will imply a different perturbation dynamics. We shall demonstrate that non-adiabatic features appear as a characteristic signature for a large class of interacting models. These effects may be used to discriminate between different models of the cosmic medium which share the same background dynamics.

The outline of this paper is as follows. Section II provides the general formalism for an interacting two-component fluid, where one of the fluids is pressureless. Then it focuses on the case for which the ratio of the energy densities of both fluids is constant or slowly varying. This will single out models for which the energy density of the second component is proportional to the square of the Hubble parameter. A realization of this dependence is provided by certain holographic dark energy models. The basic properties of these models are recalled in Section III. Section IV discusses in detail matter in interaction with holographic dark energy. It provides the conditions under which an interaction driven transition from decelerated to accelerated expansion is feasible. It also comments on the possibility of a weak time dependence of the saturation parameter of the holographic bound, which is usually assumed constant. Section V introduces a simple interacting model and compares it with the Λ CDM model. The perturbation dynamics for a fluctuating interaction rate is discussed in VI with special emphasis on non-adiabatic aspects on large perturbation scales. Finally, section VII summarizes our findings.

II. INTERACTING COSMOLOGICAL FLUIDS

The field equations for a spatially homogeneous and isotropic universe are the Friedmann equation

$$3 H^2 = 8 \pi G \rho - 3 \frac{k}{a^2}, \quad (k = +1, 0, -1), \quad (1)$$

and

$$\dot{H} = -4 \pi G (\rho + p) + \frac{k}{a^2}. \quad (2)$$

A combination of both equations yields

$$2 \frac{\dot{H}}{H} = -3H \left(1 + \frac{p}{\rho}\right) - \frac{k}{a^2 H} \left(1 + 3 \frac{p}{\rho}\right), \quad (3)$$

a relation that will be useful later on. The deceleration parameter, $q = -\ddot{a}/(aH^2)$, can be written as

$$q = \frac{1}{2} \left(1 + \frac{k}{a^2 H^2}\right) \left(1 + 3 \frac{p}{\rho}\right). \quad (4)$$

The total energy density ρ is supposed to split into $\rho = \rho_M + \rho_X$, where ρ_M is the energy density of pressureless dark matter. Under this assumption the total pressure equals the dark energy pressure, $p = p_X$. We further assume that both components do not conserve separately but interact with each other in such a manner that the balance equations take the form

$$\dot{\rho}_M + 3H\rho_M = Q \quad \text{and} \quad \dot{\rho}_X + 3H(1+w)\rho_X = -Q, \quad (5)$$

where $w \equiv p_X/\rho_X$ is the equation of state parameter of the dark energy, and the function $Q > 0$ measures the strength of the interaction. Models featuring an interaction matter–dark energy were introduced by Wetterich [34] (see also [35]) and first used alongside the holographic dark energy by Horvat [29]. Nowadays there is a growing body of literature on the subject -see, e.g. [36] and references therein. Although the assumption of a coupling between both components implies the introduction of an additional phenomenological function Q , a description that admits interactions is certainly more general than otherwise. Further, there is no known symmetry that would suppress such interaction and arguments in favor of interacting models have been put forward recently [37].

The quantity of interest for analyzing the coincidence problem is the ratio $r \equiv \rho_M/\rho_X$, which, upon using Eqs. (5), can be written as

$$\dot{r} = (1+r) \left[3Hw \frac{r}{1+r} + \Gamma \right] . \quad (6)$$

Here we have introduced the quantity $\Gamma \equiv Q/\rho_X$ which characterizes the rate by which ρ_X changes as a result of the interaction. On the other hand, combining the balance equation for ρ_X in (5) with Eq. (3), we obtain

$$\frac{\dot{\rho}_X}{\rho_X} - 2 \frac{\dot{H}}{H} = - \left[3Hw \frac{r}{1+r} + \Gamma \right] + \frac{k}{a^2 H} \left(1 + 3 \frac{w}{1+r} \right) . \quad (7)$$

Comparing now Eqs. (7) and (6), it follows that the dynamics of the aforesaid ratio is governed by

$$\dot{r} = - (1+r) \left[\frac{\dot{\rho}_X}{\rho_X} - 2 \frac{\dot{H}}{H} - \frac{k}{a^2 H} \left(1 + 3 \frac{w}{1+r} \right) \right] . \quad (8)$$

It is this formula which deserves attention with regard to the coincidence problem. Clearly, the lower $|r|$, the less acute this problem seems to be. Inspection of Eq. (8) shows that the case $\rho_X \propto H^2$ is singled out for it leads to

$$\dot{r} = (1+r) \frac{k}{a^2 H} \left(1 + 3 \frac{w}{1+r} \right) . \quad (9)$$

At this point, it is expedient to introduce the dimensionless quantities

$$\Omega_M = \frac{8\pi G \rho_M}{3 H^2}, \quad \Omega_X = \frac{8\pi G \rho_X}{3 H^2}, \quad \Omega_k = -\frac{k}{a^2 H^2}. \quad (10)$$

Thus, Friedmann's equation (1) can be cast as

$$\Omega_M + \Omega_X + \Omega_k = 1 . \quad (11)$$

By using it together with Eq. (4), the expression (9) takes the form

$$\dot{r} = -2H \frac{\Omega_k}{\Omega_X} q . \quad (12)$$

Via the curvature the evolution of r is directly linked to the deceleration parameter. For $k = +1$ and $q > 0$ (decelerated expansion) r augments, and for $q < 0$ (accelerated expansion)

r diminishes. For $k = -1$ the behavior is just the opposite. In any case, only small variations of r are possible nowadays if $1/(a^2 H^2) \ll 1$ at present [3, 16]. While a slow time dependence of the energy density ratio is desirable from the point of view of the coincidence problem, it remains to be clarified whether dark energy models with $\rho_X \propto H^2$ are able to account for a present phase of accelerated expansion as well as for a transition to the latter from an earlier matter dominated phase.

A survey of the, by now, rather ample body of literature on dark energy candidates reveals that models of this type have indeed been introduced previously, namely, in the context of ideas which are rooted in the holographic principle. In the following section we briefly recall, how a dependence $\rho_X \propto H^2$ emerges in these holographic dark energy models.

III. HOLOGRAPHIC DARK ENERGY

The basic ideas of the holographic principle were introduced by 't Hooft [25] and Susskind [26]. The development of interest for our purpose was put forward by Cohen *et al.* [27], followed by Hsu [28] and Li [31], who considered specific holographic dark energy models.

The essential point in establishing the holographic idea is the way of counting the degrees of freedom of a physical system. Consider a three-dimensional lattice of spin-like degrees of freedom and assume that the distance between every two neighboring sites is some small length ℓ which is of the order of the Planck length, ℓ_{Pl} . Each spin can be in one of two states. In a region of volume L^3 the number of quantum states N will be $N = 2^n$, with $n = (L/\ell)^3$ the number of sites in the volume, whence the entropy, given by the logarithm of N , will be $S \propto (L/\ell)^3 \ln 2$. Identifying (in Planck units) ℓ^{-1} with the ultraviolet cutoff Λ (the corresponding energy is $L^3 \Lambda^4$), the maximum entropy varies as $S \sim L^3 \Lambda^3$, i.e., proportional to the volume of the system. Based on considerations on black hole thermodynamics (bear in mind that the Bekenstein–Hawking entropy is $S_{BH} = A/(4 \ell_{Pl}^2)$, where A is the area of the black hole horizon), Bekenstein [38] argued that the maximum entropy for a box of volume L^3 should be proportional to its surface rather than to its volume. In keeping with this, 't Hooft conjectured that all phenomena within a volume L^3 should be described by a set of degrees of freedom located at the surface which bounds this volume with approximately one binary degree of freedom per Planck's area.

Inspired by these ideas, Cohen *et al.* [27] demonstrated that an effective field theory that

saturates the inequality (the Bekenstein bound)

$$L^3 \Lambda^3 \leq S_{BH} \simeq L^2 M_{Pl}^2 \quad (M_{Pl}^2 = (8\pi G)^{-1}), \quad (13)$$

necessarily includes states for which the Schwarzschild radius R_s is larger than the box size, i.e., $R_s > L$. Namely, for sufficiently high temperatures (in Planck units $T \gg L^{-1}$ but $T \leq \Lambda$) the thermal energy of the system is $E \simeq L^3 T^4$ while its entropy is $S \simeq L^3 T^3$. When Eq. (13) is saturated (by setting $T = \Lambda$ in there) it follows that $T \simeq (M_{Pl}^2/L)^{1/3}$. Now, the Schwarzschild radius R_s is related to the energy by $R_s \simeq E/M_{Pl}^2$ (recall that an object with Schwarzschild radius R_s corresponds to a (Newtonian) mass $R_s/(2G)$). Consequently, $R_s \gg L$, i.e., the Schwarzschild radius is indeed larger than the system size.

A stronger constraint which excludes states for which the Schwarzschild radius exceeds the size L is

$$L^3 \Lambda^4 \leq M_{Pl}^2 L. \quad (14)$$

The expression on the right-hand side of Eq. (14) corresponds to the energy of a black hole of size L . So, this constraint ensures that the energy $L^3 \Lambda^4$ in a box of the size L does not exceed the energy of a black hole of the same size [31].

While the Bekenstein bound (13) implies a scaling $L \propto \Lambda^{-3}$, the inequality (14) corresponds to a behavior $L \propto \Lambda^{-2}$. Furthermore, since saturation of (14) means $\Lambda^3 \simeq (M_{Pl}^2/L^2)^{3/4}$, one finds that $S = L^3 \Lambda^3 = (M_{Pl}^2 L^2)^{3/4} = S_{BH}^{3/4}$.

By saturating the inequality (14) and identifying Λ^4 with the holographic energy density ρ_X it follows [27, 31]

$$\rho_{HDE} = \frac{3c^2}{8\pi G L^2}, \quad (15)$$

where the factor 3 was introduced for convenience and c^2 is a dimensionless quantity which is usually assumed constant. It is obvious, that for a choice $L = H^{-1}$, i.e., for a cutoff set by the Hubble radius, the energy density (15) is characterized by exactly that dependence $\propto H^2$ which was singled out by Eqs. (8) and (9). In the following we shall identify (15) with the energy density ρ_X appearing in Eq. (5).

IV. HOLOGRAPHIC DARK ENERGY IN INTERACTION WITH MATTER

For holographic dark energy models with a cutoff scale H^{-1} the time dependence of the energy density ratio is determined by Eq. (9). For a spatially flat universe the ratio

r remains constant. Possible changes of r due to a non-vanishing spatial curvature term are necessarily small. At first glance, models with a constant (or almost constant) energy density ratio seem unable to account for a transition from decelerated to accelerated expansion. And indeed, holographic dark energy models with the Hubble scale as cutoff length were considered to be ruled out since their equation of state parameter does not seem to allow negative values which are required for an accelerated expansion [28, 31]. However, as shown previously, the presence of an interaction between dark energy and dark matter may change this situation [33]. This becomes obvious if Eq. (3) with $p = w\rho_X$ is solved for the equation of state parameter w . The result is

$$w = -\frac{1}{1 - \Omega_X} \left(\frac{\Gamma}{3H} + \frac{\Omega_k}{3} \right) . \quad (16)$$

Assuming $|\Omega_k|$ small, the interaction rate Γ essentially determines the equation of state parameter as soon as the dimensionless ratio Γ/H , which here and in the following is the relevant interaction parameter, becomes of order one. For negligible interaction, $|\Gamma|/H \ll 1$, one has $|w| \ll 1$.

The main point of interest now is the behavior of the deceleration parameter, i.e., whether this kind of models admits a transition from decelerated to accelerated expansion. The condition for accelerated expansion ($q < 0$)

$$w + \frac{\Gamma}{3H} < -\frac{1}{3} , \quad (17)$$

readily follows from the above equations. Introduction of the expression (16) for w into the latter leads to

$$\frac{\Gamma}{H} > \frac{1 - \Omega_X}{\Omega_X} - \frac{\Omega_k}{\Omega_X} \quad \Leftrightarrow \quad \frac{\Gamma}{H} > r , \quad (18)$$

where we have used Friedmann's equation (11). On the other hand, since holographic energy is not compatible with phantom fields [39], one must impose $w \geq -1$ to be consistent with the holographic idea. From Eq. (16) we find that this condition amounts to

$$\frac{\Gamma}{H} \leq 3 (1 - \Omega_X) - \Omega_k . \quad (19)$$

Therefore, the consistency condition simply reads,

$$3\Omega_X > 1 - \Omega_k . \quad (20)$$

The inequality (19) says that the ratio on its left hand side is smaller than a quantity of order one. Clearly, this is compatible with an initial condition $|\Gamma|/(3H) \ll 1$. The latter implies that w must be close to zero deep in the matter dominated era (cf. Eq. (16)). An initial condition $|\Gamma| \ll H$ is equivalent to an initially negligible interaction. This means that at high redshifts (but well after matter–radiation equality), the dark energy component is almost non–interacting and its equation of state is close to that of pressureless matter. On the other hand, the condition (18) for accelerated expansion entails that $|\Gamma|/H$ must be larger than a lower bound of order one, as well. Both inequalities are consistent with condition (20). This supports our suggestion that, depending on the interaction rate, a transition from decelerated to accelerated expansion is indeed feasible.

Starting from Eq. (8), the dynamics of the energy density ratio may alternatively be written as

$$\dot{r} = H \frac{\Omega_k}{1 - \Omega_X} \left[\frac{\Gamma}{H} - r \right]. \quad (21)$$

For $k = 0$ we recover the stationary case $r = \text{constant}$, irrespective of the value of Γ/H . If additionally $\Gamma/H = \text{constant}$, it follows from Eq. (16) that w is also a constant. In such a case, there is no transition from decelerated to accelerated expansion [33]. However, if Γ/H is allowed to grow, then the transition may well occur, even though the ratio r remains constant or almost constant (see Eq. (24) below).

The spatially curved cases generate an interesting additional dynamics. The transition condition $\Gamma/H = r$ (cf. Eq. (18)) coincides with the condition for $\dot{r} = 0$ for $k = \pm 1$ in (21). Since at early times one has $\Gamma/H \ll r$, the ratio r in (21) increases for $k = +1$, while it decreases for $k = -1$. It reaches a maximum (minimum) for $k = 1$ ($k = -1$) at $\Gamma/H = r$ and may go down (up) afterward.

The obvious requirement for a sufficient growth of the interaction parameter Γ/H is that $(\Gamma/Hr)' > 0$ in the phase of decelerated expansion or, by (21), that

$$\frac{(\Gamma/H)'}{\Gamma/H} > -\frac{H \Omega_k}{1 - \Omega_X} \left[1 - \frac{1}{r} \frac{\Gamma}{H} \right]. \quad (22)$$

But according to Eq. (19) the ratio Γ/H is bounded from above. For a growth of Γ/H from a very small value to a value of order one, a transition from decelerated to accelerated expansion is feasible. We mention again, that any change of r arises solely as a curvature

effect.

A different way to understand the dynamics of the ratio r , which also may be seen as a consistency check, is to start from Eq. (11) and to realize that $\Omega_X = \text{constant}$. Then (recall that $r = \Omega_M/\Omega_X$),

$$r = \frac{1 - \Omega_X}{\Omega_X} - \frac{\Omega_k}{\Omega_X}. \quad (23)$$

The first term on the right hand side of (23) is constant. It is just the curvature term that makes r vary. Since in accelerating universes $|\Omega_k|$ decreases r will approach a constant value in the long time limit. The ratio r grows if the curvature term enlarges the right hand side ($k = 1$ and $q > 0$, or $k = -1$ and $q < 0$) and it goes down if the curvature term diminishes the right hand side ($k = 1$ and $q < 0$, or $k = -1$ and $q > 0$). This is essentially the behavior discussed beneath Eq. (12). Clearly, differentiation of Eq. (23) alongside the introduction of the deceleration parameter q consistently reproduces Eq. (12).

Combining Eqs. (12) and (21) we get

$$\frac{\Gamma}{H} - r = -2 \frac{1 - \Omega_X}{\Omega_X} q. \quad (24)$$

This equation encodes the central message of our paper. It covers all the cases $k = 0, \pm 1$ and explicitly shows that the cosmological dynamics crucially depends on the difference of two ratios: the ratio Γ/H and the ratio r of the energy densities. The difference of these two ratios is directly proportional to the (negative) deceleration parameter. Accelerated expansion requires $\Gamma/H > r$, decelerated expansion demands $\Gamma/H < r$. It is obvious, that even a constant or a slowly varying ratio r may be compatible with a sign change of q provided Γ/H evolves accordingly. Even with a constant r a present accelerated expansion is compatible with an earlier matter dominated period with decelerated expansion. The point is that for negligible interaction (at sufficiently high redshifts) the dark energy component behaves as non-relativistic matter. In this phase r is not really an important parameter since it describes the ratio of two components with the same equation of state. It was this property that apparently ruled out a (non-interacting) holographic dark energy model with an infrared cutoff set by the Hubble scale [28, 31]. Here, this unwanted (in the non-interacting model) feature is advantageous since, thanks to it, a matter dominated phase during which structure formation can occur is naturally recovered. It is only because of the gradually increased interaction that the equations of state begin to differ from each other. Then r becomes important. At the first glance the circumstance that r is constant or only

slowly varying seems to imply, that the coincidence problem gets significantly alleviated. But in fact, it has been shifted from the problem to explain a constant (or slowly varying) ratio r towards the problem to explain an interaction rate which has to be of the order of the Hubble rate just at the present time. We argue that an alternative, dynamical formulation of the coincidence problem maybe useful, in particular, if a specific interacting model of the type presented here will lead to potentially observable differences from, say the Λ CDM model or generalized Chaplygin gas models (see below).

One might argue against a constant -or nearly constant- ratio r by saying that, as is well known, at the time of primeval nucleosynthesis the dark energy should not contribute more than 5 per cent to the total energy density if the standard big bang scenario for the build up of light elements is to hold [41]. However, this criticism does not apply to our case since, as said above, the equation of state of dark energy, w , was not close to -1 at early epochs but close to that of dust.

By starting from the expression (15) with $L = H^{-1}$ for the dark energy we have assumed the parameter c^2 to be constant throughout our considerations. However, there does not seem to exist any compelling reason for the inequality (14) to be saturated or for the degree of saturation to be a constant once and for all. In principle, a weak time dependence ($0 < (c^2)' / c^2 \ll H$) of c^2 should be admitted here. Such an additional degree of freedom will modify the dynamics discussed so far. In particular, instead of described by Eq. (21), the ratio r will evolve according to

$$\dot{r} = -H r \frac{1}{1 - \Omega_X} \left\{ -\Omega_k \left[\frac{1}{r} \frac{\Gamma}{H} - 1 \right] + \frac{(c^2)'}{H c^2} \right\}, \quad (25)$$

and the equation of state parameter changes from (16) to

$$w = -\frac{1}{1 - \Omega_X} \left[\frac{\Gamma}{3H} + \frac{\Omega_k}{3} + \frac{1}{3H} \frac{(c^2)'}{c^2} \right]. \quad (26)$$

As a consequence, in the conditions (17) and (18) and in all relations after them, the rate Γ is to be replaced by $\Gamma + \frac{(c^2)'}{c^2}$. A varying c^2 will induce small variations in r , additionally to those which are due to the spatial curvature and can support the transition from decelerated to accelerated expansion [33] (see also [40]). Thus, at this transition, we have

$$\dot{r}(q=0) = -\frac{r}{1-\Omega_X} \frac{(c^2)^\cdot}{c^2} \left(1 + \frac{\Omega_k}{r}\right), \quad (27)$$

which reduces to the previous result $\dot{r}(q=0) = 0$ for $c^2 = \text{constant}$.

Before closing this section, it is noteworthy that a negative equation of state parameter w can be obtained from a negative curvature term ($k = -1$) alone (cf. Eq. (16)), i.e., even for $\Gamma = 0$,

$$w = -\frac{1}{3} \frac{1}{a^2 H^2} \frac{1}{1-\Omega_X}. \quad (28)$$

However, the condition (18) for accelerated expansion, in this case $1/(aH)^2 > 1 - \Omega_X$, can never be satisfied as it contradicts the constraint equation (11). Consequently, only models with a non-vanishing interaction between dark energy and dark matter can be potential candidates for a satisfactory cosmological dynamics based on the holographic idea with the infrared cutoff set by the Hubble function.

V. A SIMPLE MODEL

In our approach, the unknown nature of dark energy is mapped on a (so far) unspecified interaction in the dark sector quantified by the ratio Γ/H . In this section we build a simple model that explicitly exhibits the general features discussed in the preceding section.

With the help of the equation of state parameter (16) the balances expressions (5) may be written as

$$\frac{\dot{\rho}_X}{\rho_X} = -3H \left[1 - \frac{1}{3} \frac{\Omega_k}{1-\Omega_X} - \frac{\Gamma}{3H} \frac{\Omega_X}{1-\Omega_X} \right], \quad \frac{\dot{\rho}_M}{\rho_M} = -3H \left[1 - \frac{\Gamma}{3H} \frac{\Omega_X}{\Omega_M} \right]. \quad (29)$$

For $k = 0$, we have $1 - \Omega_X = \Omega_M$ and both rates coincide. Moreover, in this case r stays constant. Integration of (29.2) under this condition from some initial period (subscript “ i ”) onward, results in

$$\frac{\rho_M}{\rho_{Mi}} = \left(\frac{a_i}{a}\right)^3 \exp \left[\frac{1}{r} \int \frac{da}{a} \frac{\Gamma}{H} \right]. \quad (30)$$

The exponential describes the deviation from an Einstein - de Sitter universe. For the special case of a constant Γ/H , the expression (30) reduces to

$$\frac{\rho_M}{\rho_{Mi}} = \left(\frac{a_i}{a}\right)^{3-\Gamma/(rH)} . \quad (31)$$

Consequently, the Hubble parameter depends on the scale factor as $H \propto a^{-\frac{1}{2}[3-\Gamma/(rH)]}$. Hence,

$$a \propto t^{\frac{2}{3-\Gamma/(rH)}} . \quad (32)$$

The exponent on the right hand side of the last expression is larger than unity for $\Gamma/(rH) > 1$ which reproduces the previous condition (18) for accelerated expansion. The limiting case $\Gamma/(rH) = 3$ corresponds to a constant energy density, i.e., to an exponential growth of the scale factor.

As mentioned above, a constant ratio Γ/H can describe either a phase of decelerated expansion ($\Gamma/(rH) < 1$), or a phase of accelerated expansion ($\Gamma/(rH) > 1$), but never a transition between the two. To stage a transition from decelerated to accelerated expansion $\Gamma/(rH)$ must increase which, for a constant ratio r , means that Γ/H must increase. Since there is no guidance of what a microscopic interaction model could be, we shall resort to a phenomenological approach, assuming a growth of the ratio Γ/H with a power of the scale factor a . Notice that a working microscopic interaction model in the present context, where it is exclusively the interaction that drives the accelerated expansion, would be equivalent to explaining the nature of dark energy. The fact that we obtain a transition from decelerated to accelerated expansion under the condition of a constant energy density ratio makes the coincidence problem appear in a different way. In our approach it remains to explain why the interaction rate is of the order of the Hubble rate just at the present time. We argue, that this reformulation of the coincidence problem may be advantageous since it offers a potential dynamical solution, albeit hypothetical at the present state of knowledge.

In the following we assume a growth according to

$$\frac{\Gamma}{rH} = 3\beta \left(\frac{a}{a_0}\right)^\alpha , \quad a \leq a_0, \quad \beta \lesssim 1 , \quad (33)$$

with constant, positive-definite parameters α and β . The condition on β ensures, that the phantom divide cannot be crossed. Furthermore, we restrict our considerations to the past evolution of the universe. The ansatz (33) implies a continuous growth of the ratio $\Gamma/(rH)$ with a maximum value at the present time. We do not speculate about a possible influence of the interaction on the future dynamics of the Universe. But it is obvious that if, in the future, the interaction becomes less effective, an evolution back to a fresh phase of decelerated expansion is feasible.

Inserting the ansatz (33) into Eq. (30) we get

$$\frac{\rho_M}{\rho_{Mi}} = \left(\frac{a_i}{a}\right)^3 \exp \left[\frac{3\beta}{\alpha} \left(\left(\frac{a}{a_0}\right)^\alpha - \left(\frac{a_i}{a_0}\right)^\alpha \right) \right]. \quad (34)$$

The exponential factor diminishes the decreases of the energy density with cosmic expansion. Replacing the initial quantities by the corresponding present day values (subscript “0”) provides us with

$$\rho_M = \rho_{M0} \left(\frac{a_0}{a}\right)^3 \exp \left[\frac{3\beta}{\alpha} \left(\left(\frac{a}{a_0}\right)^\alpha - 1 \right) \right]. \quad (35)$$

For $k = 0$ and $c^2 = \text{constant}$, the energy density of the dark energy component shows exactly the same dependence on the scale factor,

$$\rho_X = \rho_{X0} \left(\frac{a_0}{a}\right)^3 \exp \left[\frac{3\beta}{\alpha} \left(\left(\frac{a}{a_0}\right)^\alpha - 1 \right) \right]. \quad (36)$$

Thus, the Hubble rate is given by

$$h \equiv \frac{H}{H_0} = \left(\frac{a_0}{a}\right)^{3/2} \exp \left[\frac{3\beta}{2\alpha} \left(\left(\frac{a}{a_0}\right)^\alpha - 1 \right) \right]. \quad (37)$$

Comparing Eq. (33) with (24) for the spatially flat case, the deceleration parameter is found to be

$$q = \frac{1}{2} - \frac{3}{2} \beta \left(\frac{a}{a_0}\right)^\alpha. \quad (38)$$

It is expedient to compare the dimensionless Hubble rate, h , given by Eq. (37), with the corresponding quantity of the Λ CDM model,

$$\frac{H_{\Lambda CDM}}{H_0} = \sqrt{\frac{\rho_\Lambda}{\rho_\Lambda + \rho_{M0}}} \left[1 + \frac{\rho_{M0}}{\rho_\Lambda} \left(\frac{a_0}{a} \right)^3 \right]^{1/2}. \quad (39)$$

Introducing the redshift parameter z by $a_0/a = 1 + z$, we have

$$\begin{aligned} \frac{H_{\Lambda CDM}}{H_0} &= \sqrt{\frac{\rho_\Lambda}{\rho_\Lambda + \rho_{M0}}} \left[1 + \frac{\rho_{M0}}{\rho_\Lambda} (1 + z)^3 \right]^{1/2} \\ &= 1 + \frac{3}{2} \frac{\rho_{M0}}{\rho_\Lambda + \rho_{M0}} z + \mathcal{O}(z^2), \end{aligned} \quad (40)$$

where for reasons explained below we have retained only the linear term in z in the second line of last equation. From (37) it follows that




FIG. 1: Graphs of the dimensionless ratio h vs redshift. Except for the middle graph -which corresponds to the Λ CDM model-, from top to bottom the α values are: 2.0, 1.5, 1.2, and 1.0. Here, we have taken $\beta = 3/4$.

$$\begin{aligned}\frac{H}{H_0} &= (1+z)^{3/2} \exp \left[\frac{3\beta}{2\alpha} \left(\left(\frac{1}{1+z} \right)^\alpha - 1 \right) \right] \\ &= 1 + \frac{3}{2}(1-\beta)z + \mathcal{O}(z^2),\end{aligned}\tag{41}$$

for our interacting model. It is remarkable that to linear order in z , the expression (41) does not depend on α .

We may now require that our model (41) coincides with the Λ CDM model (40) up to linear order in z . This allows us to fix the parameter β in terms of the observationally well established ratio of the parameters ρ_{M0} and ρ_Λ of the Λ CDM model. The result is

$$\beta = \frac{\rho_\Lambda}{\rho_\Lambda + \rho_{M0}}.\tag{42}$$

For the observed $\rho_{M0}/\rho_\Lambda \approx 1/3$ we have $\beta \approx 3/4$. Under this condition, the present value of the deceleration parameter (38), $q_0 = \frac{1}{2} - \frac{3}{2}\beta$, coincides with the corresponding value $q_{\Lambda CDM0} = 1/2 - [3\rho_\Lambda/2(\rho_\Lambda + \rho_{M0})] \approx -0.6$ of the Λ CDM model as well. However, the evolution of q towards this value depends on α which gives rise to differences in any order beyond the linear one. For the value of the transition redshift z_{acc} , that follows from $q = 0$ in (38), we obtain $z_{acc} = (3\beta)^{1/\alpha} - 1$.

With the identification (42) and $\rho_{M0}/\rho_\Lambda \approx 1/3$ we find $z_{acc} \approx 1.2$ for $\alpha = 1$ and $z_{acc} \approx 0.5$ for $\alpha = 2$. The corresponding z -value for the Λ CDM model is approximately 0.8.

Figure 1 contrasts the Hubble parameter prediction of our interaction model for different values of α with the Λ CDM model. Clearly, the free parameter α , obviously of the order of one, can be used to adjust $h(z)$ according to the observational situation.

Figure 2 presents the best fit of our model and the best fit of the Λ CDM model to the Gold SNIa data set of Riess *et al.* (fourth reference in [4]), and the SNLS data set of Astier *et al.* (sixth reference in [4]). In plotting the graphs the distance modulus, $\mu = 5 \log d_L + 25$, was employed. In this expression $d_L = (1+z) \int_0^z H^{-1}(z') dz'$ is the luminosity distance in megaparsecs. As can be seen, both best fits largely overlap one another. In all, current SNIa data are unable to discriminate between the popular Λ CDM and our interaction model.




FIG. 2: Distance modulus vs redshift for the best fit model, $\Omega_X = 0.75$, $H_0 = 67.35$ km/s/Mpc, $\alpha = 1.5$ (solid line), and the Λ CDM model, $\Omega_\Lambda = 0.71$, and $H_0 = 67.35$ km/s/Mpc (dashed line).

VI. PERTURBATION DYNAMICS

In a flat universe the dark energy equation of state parameter is given by $w = -(1+r)\frac{\Gamma}{3Hr}$. Since $\rho_X = \frac{\rho}{1+r}$, the total equation of state of the cosmic medium is

$$\frac{p}{\rho} = -\frac{\Gamma}{3Hr}. \quad (43)$$

The corresponding adiabatic sound speed parameter becomes

$$\frac{\dot{p}}{\dot{\rho}} = \left(1 - \frac{1}{3\gamma H} \frac{(\Gamma/H)^\cdot}{\Gamma/H}\right) \frac{p}{\rho}, \quad \left(\gamma = 1 + \frac{p}{\rho}\right). \quad (44)$$

However, the interaction introduces non-adiabatic features into the perturbation dynamics. Deviations from the adiabatic behavior are suitably characterized by the quantity $\hat{p} - \frac{\dot{p}}{\dot{\rho}}\hat{\rho}$, which in our case takes the form

$$\hat{p} - \frac{\dot{p}}{\dot{\rho}}\hat{\rho} = p \left[\frac{(\Gamma/H)^{\hat{}}}{\Gamma/H} - \frac{(\Gamma/H)^{\cdot}}{\Gamma/H} \frac{\hat{\rho}}{\dot{\rho}} - \frac{\hat{r}}{r} \right] , \quad (45)$$

where a hat indicates perturbation of the corresponding quantity. The contribution on the right-hand side of (45) accounts both for the non-adiabaticity of the X component and for a contribution which results from the two component nature of the system. In terms of the components this expression is equivalent to

$$P - \frac{\dot{p}}{\dot{\rho}}D = \frac{\dot{\rho}_X}{\dot{\rho}} \left(P_X - \frac{\dot{p}_X}{\dot{\rho}_X} D_X \right) + \frac{\dot{\rho}_M \dot{\rho}_X}{\dot{\rho}^2} \frac{\dot{p}_X}{\dot{\rho}_X} [D_X - D_M] , \quad (46)$$

where

$$D \equiv \frac{\hat{\rho}}{\rho + p} = -3H \frac{\hat{\rho}}{\dot{\rho}} , \quad (47)$$

and the energy density perturbations for the components are

$$D_M \equiv -3H \frac{\hat{\rho}_M}{\dot{\rho}_M} \quad \text{and} \quad D_X \equiv -3H \frac{\hat{\rho}_X}{\dot{\rho}_X} . \quad (48)$$

Analogously, the pressure perturbations are described by

$$P \equiv \frac{\hat{p}}{\rho + p} = -3H \frac{\hat{p}}{\dot{\rho}} \quad \text{and} \quad P_X \equiv -3H \frac{\hat{p}_X}{\dot{\rho}_X} . \quad (49)$$

The circumstance that $\dot{r} = 0$ in these class of models results in a particularly simple structure for $D_X - D_M$. Just from the definition of these quantities we find by direct calculation

$$D_X - D_M = -\frac{\hat{r}}{\gamma r} . \quad (50)$$

This difference is directly related to the fluctuation of the energy density ratio. The dynamic equation for $D_X - D_M$ (see below) then determines the time dependence of $\hat{r}$.

A homogenous and isotropic background universe with scalar metric perturbations and vanishing anisotropic pressure can be characterized by the line element (longitudinal gauge, cf. [42])

$$ds^2 = -(1 + 2\psi) dt^2 + a^2 (1 - 2\psi) \delta_{\alpha\beta} dx^\alpha dx^\beta . \quad (51)$$

The perturbation dynamics is most conveniently described in terms of the gauge-invariant variable [43]

$$\zeta \equiv -\psi + \frac{1}{3} \frac{\hat{\rho}}{\rho + p} = -\psi - H \frac{\hat{\rho}}{\dot{\rho}} , \quad (52)$$

which represents curvature perturbations on hypersurfaces of constant energy density. Corresponding quantities for the components are

$$\zeta_A \equiv -\psi - H \frac{\hat{\rho}_A}{\dot{\rho}_A} \quad (A = X, M). \quad (53)$$

On large perturbation scales the variable ζ obeys the equation (cf. [44, 45, 46])

$$\dot{\zeta} = -H \left(P - \frac{\dot{p}}{\dot{\rho}} D \right). \quad (54)$$

The equation for $\zeta_X - \zeta_M = \frac{1}{3} (D_X - D_M)$ is

$$\begin{aligned} (\zeta_X - \zeta_M)^{\cdot} = & 3H^2 \frac{\hat{p}_X - \frac{\dot{p}_X}{\dot{\rho}_X} \hat{\rho}_X}{\dot{\rho}_X} \\ & + 3H \frac{\dot{\Pi}}{\dot{\rho}_M \dot{\rho}_X} \left[\zeta_{\pi} - \zeta + \frac{\dot{\rho}_X - \dot{\rho}_M}{\dot{\rho}} (\zeta_X - \zeta_M) \right]. \end{aligned} \quad (55)$$

The gauge invariant quantity ζ_{Π} describes the perturbation of the interaction term. It is defined in analogy to the other perturbation quantities,

$$\zeta_{\Pi} \equiv -\psi - H \frac{\hat{\Pi}}{\dot{\Pi}}, \quad \Pi = -\frac{\Gamma \rho_X}{3H}. \quad (56)$$

Inserting in (55) the expression (50) and

$$\hat{p}_X - \frac{\dot{p}_X}{\dot{\rho}_X} \hat{\rho}_X = -\frac{\Gamma \rho}{3Hr} \left[\frac{(\Gamma/H)^{\cdot}}{\Gamma/H} - \frac{\hat{r}}{r(1+r)} + \frac{1}{3\gamma H} \frac{(\Gamma/H)^{\cdot}}{\Gamma/H} \left(\frac{\hat{\rho}}{\rho} - \frac{\hat{r}}{1+r} \right) \right], \quad (57)$$

which follows from the equation of state (43), we obtain

$$\dot{\hat{r}} = 0 \quad \Rightarrow \quad \hat{r} = \text{const} \quad (58)$$

on large perturbation scales, i.e. scales for which spatial gradient terms may be neglected. This means, also at the perturbative level, $\hat{r}$ is not a dynamical degree of freedom on these scales. It is remarkable that this property holds for any interaction rate. While, of course, the trivial case $\hat{r} = 0$ is included here, it is expedient to notice that any non-vanishing $\hat{r}$, although constant, gives rise to non-adiabatic effects.

The simplest case to discuss this dynamics more specifically is $\Gamma = \text{const}$. Since $H \propto \rho^{1/2}$, this corresponds to an effective background equation of state $p \propto -\rho^{1/2}$ which is

characteristic for a special generalized Chaplygin gas [47]. Its energy density is

$$\rho = \rho_i \left[\frac{\Gamma}{3H_i r} + \left(1 - \frac{\Gamma}{3H_i r} \right) \left(\frac{a_i}{a} \right)^{3/2} \right]^2 . \quad (59)$$

Again, for $\Gamma = 0$ the Einstein de Sitter universe is reproduced. The general relations (44) and (45) specify to

$$\frac{\dot{p}}{\dot{\rho}} = \frac{1}{2} \frac{p}{\rho} \quad (60)$$

and

$$\hat{p} - \frac{\dot{p}}{\dot{\rho}} \hat{\rho} = \frac{1}{2} p \left[\frac{\hat{\rho}}{\rho} - 2 \frac{\hat{\Theta}}{\Theta} - \frac{\hat{r}}{r} \right] \quad (\Theta = 3H) , \quad (61)$$

respectively. Only if spatial gradient terms in the perturbation dynamics are neglected, the first two terms in the bracket on the right-hand side of Eq. (61) cancel each other approximately. Therefore one may expect that non-adiabatic effects will be most important on small perturbation scales [48, 49]. But even for $\frac{\hat{\rho}}{\rho} \approx 2 \frac{\hat{\Theta}}{\Theta}$ a non-vanishing perturbation $\hat{r}$ generates a non-adiabatic dynamics of the system. The corresponding large scale perturbation dynamics can be solved analytically. The equation for ζ in this case reduces to

$$\dot{\zeta} = - \frac{\Gamma}{6\gamma} \frac{\hat{r}}{r^2} . \quad (62)$$

With $\gamma = 1 + \frac{p}{\rho} = 1 - \frac{\Gamma}{3Hr}$ we may write

$$\zeta' = - \frac{\Gamma}{2} \frac{\hat{r}}{r} \frac{1}{a} \frac{1}{3Hr - \Gamma} , \quad (63)$$

where $\zeta' \equiv \frac{d\zeta}{da}$. Since (cf. (59))

$$H = H_i \left[\frac{\Gamma}{3H_i r} + \left(1 - \frac{\Gamma}{3H_i r} \right) \left(\frac{a_i}{a} \right)^{3/2} \right] , \quad (64)$$

equation (63) is readily solved. The result is

$$\zeta = \zeta_i - \frac{\Gamma}{3} \frac{\hat{r}}{r} \frac{1}{3H_i r - \Gamma} \left[\left(\frac{a}{a_i} \right)^{3/2} - 1 \right] . \quad (65)$$

The non-adiabatic contribution grows with $a^{3/2}$. This will influence the time dependence of the gravitational potential and hence the integrated Sachs-Wolfe effect (cf [53, 54] for similar studies). Recall that the Λ CDM model is characterized by a constant value of ζ . On small scales we expect a modification of the adiabatic sound speed [48, 49]. Notice that the quantity $\frac{\dot{p}}{\dot{\rho}}$ in (60) is negative. As a consequence, there occur small scale instabilities

within an adiabatic perturbation analysis of this generalized Chaplygin gas. This apparently unrealistic feature has been used as an argument to discard this type of models altogether [50]. On the other hand, it has been suggested that non-adiabatic effects might modify this picture [51, 52, 53]. That this is indeed the case, has been demonstrated recently by an explicit numerical analysis for the galaxy power spectrum [49]. The present consideration in the context of interacting holographic models offer a systematic procedure to include such effects. We plan to perform a more quantitative non-adiabatic perturbation analysis in a subsequent work.

VII. DISCUSSION

We have established a holographic dark energy model in which a negative equation of state parameter as well as the transition from decelerated to accelerated expansion arise from pure interaction phenomena. A remarkable property of the model is the fact that, in a spatially flat universe, the ratio of the energy densities of dark matter and dark energy remains constant during the transition. The non-interacting limit of this model is an Einstein-de Sitter universe which is supposed to describe our cosmos at high redshifts. At first sight, the fact that r was never large may seem at variance with the conventional scenario of cosmic structure formation as one may think that at early times the amount of dark matter may have been insufficient to produce gravitational potential wells deep enough to lead to the condensation of the galaxies. However, this is not so; an earlier matter dominated phase is naturally recovered since for negligible interaction at high redshifts the equation of state of the dark energy is similar to that of non-relativistic matter. In the context of our approach the coincidence problem is dynamized and takes the form: “why is the interaction rate that drives the accelerated expansion of the order of the Hubble rate precisely at the present epoch?” Based on the assumption that the relevant interaction parameter grows with a power law in the scale factor (Eq. (33)), we have worked out a specific model for the transition from decelerated to accelerated expansion under the condition of a constant ratio of the energy densities of dark matter and dark energy. A preliminary analysis shows that this model fits the SNIa data not less well than the Λ CDM model.

We have also shown that taking into account the spatial curvature term gives rise to an additional dynamics which implies a small (compared with the Hubble rate) change of the

energy density ratio.

Furthermore, we discussed to what extent a slowly varying saturation parameter of the holographic bound may modify the cosmological dynamics. Besides, our approach can cope with a later transition to a new decelerated phase of expansion [55]-something incompatible with holographic models whose infrared cutoff is set by the radius of the future event horizon.

The interaction between dark matter and dark energy introduces non-adiabatic features into the perturbation theory. For the special case of a constant decay rate of the dark energy, in the background equivalent to a generalized Chaplygin gas, we find, Eq. (65), that the large scale curvature perturbations on hypersurfaces of constant density (which are constant in the adiabatic case, in particular for the Λ CDM model) vary with the power $3/2$ of the scale factor.

Acknowledgments

W.Z. was supported by a Grant from the “Programa de Movilidad de Profesores Visitantes 2006” sponsored by the “Direcció General de Recerca de Catalunya”. He also acknowledges support by the Brazilian grants 308837/2005-3 (CNPq) and 093/2007 (CNPq and FAPES). This work was partially supported by the Spanish Ministry of Education and Science under Grant FIS 2006-12296-C02-01, the “Direcció General de Recerca de Catalunya” under Grant 2005 SGR 000 87.

-
- [1] M. S. Turner, in *Generation of Cosmological Large-Scale Structure*, (edited by D. N. Schramm and P. Galeotti, Kluwer Academic Publishers, Dordrecht, 1997), p. 153, astro-ph/9704062.
 - [2] W. Priester *et al.*, in *Friedmann-Lemaitre Model Derived from the Lyman Alpha Forest in Quasar Spectra*, Proceedings of the Third Alexander Friedmann International Seminar on Gravitation and Cosmology, edited by Y.N. Gnedin, A.A. Grib and V.M. Mostepanenko (Friedmann Laboratory Publishing, 1995), pp 52-67.
 - [3] D. N. Spergel *et al.*, *Astrophys. J. Suppl.* **170**, 377 (2007).
 - [4] A.G. Riess *et al.*, *Astron. J.* **116**, 1009 (1998); S. Perlmutter *et al.*, *Astrophys. J.* **517**, 565 (1999); J.L. Tonry *et al.*, *Astrophys. J.* **594**, 1 (2003); A.G. Riess *et al.*, *Astrophys. J.* **607**,

- 665 (2004); M.V. John, *Astrophys. J.* **614**, 1 (2004); P. Astier *et al.*, *J. Astron. Astrophys.* **447**, 31 (2006); A.G. Riess *et al.*, astro-ph/0611572.
- [5] S. Hanany *et al.*, *Astrophys. J. Lett.* **545**, L5 2000; C.B. Netterfield *et al.*, *Astrophys. J.* **571**, 604 (2002); D.N. Spergel *et al.*, *Astrophys. J. Suppl.* **148**, 175 (2003).
- [6] M. Colless *et al.*, *Mon. Not. R. Astron. Soc.* **328**, 1039 (2001); M. Tegmark *et al.*, *Phys. Rev. D* **69**, 103501 (2004); S. Cole *et al.*, *Mon. Not. R. Astron. Soc.* **362**, 505 (2005); V. Springel, C.S. Frenk, and S.M.D. White, *Nature (London)* **440**, 1137 (2006).
- [7] S. Boughn and R. Chritenden, *Nature (London)* **427**, 45 (2004); P. Vielva, E. Martínez-González, and M. Tucci, *Mon. Not. R. Astron. Soc.* **365**, 891 (2006).
- [8] C.R. Contaldi, H. Hoekstra, and A. Lewis, *Phys. Rev. Lett.* **90**, 221303 (2003).
- [9] S. Carroll, in *Measuring and Modelling the Universe*, Carnegie Observatory, Astrophysics Series, Vol. 2, edited by W.L.Freedman, Cambridge University Press, Cambridge (2004); T. Padmanabhan, *Phys. Reports*, **380**, 235 (2003); V. Sahni, astro-ph/0403324; L. Perivolaropoulos, talk at the Third Aegean Summer School “The Invisible Universe, Dark Matter and Dark Energy”, astro-ph/0601014; E.J. Copeland, M. Sami and S. Tsujikawa, *Int. J. Mod. Phys. D* **15**, 1753 (2006).
- [10] S. Das, P.S. Corasaniti, and J. Khoury, *Phys. Rev. D* **73**, 083509 (2006).
- [11] L. Amendola, M. Gasperini, and F. Piazza, *Phys. Rev. D* **74**, 127302 (2006).
- [12] G. Olivares, F. Atrio-Barandela, and D. Pavón, *Phys. Rev. D* **71**, 063523 (2005).
- [13] G. Olivares, F. Atrio-Barandela, and D. Pavón, *Phys. Rev. D* **74**, 043521 (2006).
- [14] Zong-Kuan Guo, N. Ohta and S. Tsujikawa, *Phys. Rev. D* **76**, 023508 (2007).
- [15] W. Zimdahl and D. Pavón, *Gen. Relativ. Grav.* **36**, 1483 (2004).
- [16] U. Seljak, A. Slosar and P. McDonald, *JCAP* 10(2006)014.
- [17] R.R. Caldwell and M. Kamionkowski, *JCAP* 09(2004)009.
- [18] Bin Wang, Yungui-Gong and Ru-Keng Su, *Phys. Lett. B* **605**, 9 (2005).
- [19] J.L. Crooks, J.O. Dunn, P.H. Frampton, H.R. Norton and T. Takahashi, *Astropart. Phys.* **20**, 361 (2003), astro-ph/0305495; D. Polarski and A. Ranquet, *Phys. Lett. B* **627**, 1 (2005); E. Linder, *Astropart. Phys.* **24**, 391 (2005); K. Ichikawa and T. Takahashi, *Phys. Rev. D* **73** 083526 (2006); Zhuo-Yi Huang, Bin Wang and Ru-Keng Su, astro-ph/0605392; K. Ichikawa, M. Kawasaki, T. Sekiguchi, and T. Takahashi, *JCAP* 12(2006)005.
- [20] Yungui Gong, Bin Wang, and Yuan-Zhong Zhang, *Phys. Rev. D* **72** 043510 (2005); Qing-

- Guo Huang and Miao Li, JCAP 08(2004)013; Zhuo-Yi Huang, Bin Wang, E. Abdalla, and Ru-Keng Su, JCAP 05(2006)013.
- [21] K. Ichikawa and T. Takahashi, JCAP02(2007)001.
 - [22] Ch. Clarkson, M. Cortés and B. Bassett, JCAP 08(2007)011.
 - [23] A.A. Coley, N. Pelavas, and R.M. Zalaletdinov, Phys. Rev. Lett. **95**, 151102 (2005).
 - [24] A.A. Coley, arXiv: 0704.1734.
 - [25] G. 't Hooft, gr-qc/9310026.
 - [26] L. Susskind, J. Math. Phys. (N.Y.) **36**, 6377 (1995).
 - [27] A. G. Cohen, D.B. Kaplan and A.E. Nelson, Phys. Rev. Lett. **82**, 4971 (1999).
 - [28] S. D. H. Hsu, Phys. Lett. B **594**, 13 (2004).
 - [29] R. Horvat, Phys. Rev. D **70**, 087301 (2004).
 - [30] B. Gubina, R. Horvat and H. Nikolić, Phys. Rev. D **72**, 125011 (2005).
 - [31] M. Li, Phys. Lett. B **603**, 1 (2004).
 - [32] Yungui Gong, Phys. Rev. D **70**, 064029 (2004); Qin-Guo Huang and Yungui Gong, JCAP 08(2004)006; Bin Wang, Yungui Gong and E. Abdalla, Phys. Lett. B **624**, 141 (2005); Xin Zhang, Int. J. Mod.Phys. D **14**, 1597 (2005); Qing-Guo Huang and M. Li, JCAP 03(2005)001; Bo Hu and Yi Ling, Phys. Rev. D **73**, 123510 (2006); S. Nojiri and S.D. Odintsov, hep-th/0506212; E. Elizalde, S. Nojiri, S.D. Odintsov and Peng Wang, Phys. Rev. D **71**, 103504 (2005); Xin Zhang, Phys. Lett. B **648**, 1 (2007).
 - [33] D. Pavón and W. Zimdahl, Phys. Lett. B **628**, 206 (2005).
 - [34] C. Wetterich, Nucl. Phys. B **302**, 668 (1988); *ibid.*, Astron. Astrophys. **301**, 321 (1995).
 - [35] A. P. Billyard and A.A. Coley, Phys. Rev. D **61**, 083503 (2000).
 - [36] L. Amendola, Phys. Rev. D **62**, 043511 (2000); W. Zimdahl, D. Pavón and L.P. Chimento, Phys. Lett. B **521**, 133 (2001); L.P. Chimento, A.S. Jakubi, D. Pavón, and W. Zimdahl, Phys. Rev. D **67**, 083513 (2003); L. Amendola, Phys. Rev. D **69**, 103524 (2004); Rong-Gen Cai and Anzhong Wang JCAP 03(2005)002; Zong-Kuan Guo, Rong-Gen Cai and Yan-Zhong Zhang, JCAP 05(2005)002; W. Zimdahl, Int. J. Mod. Phys. D **14**, 2319 (2005); L.P. Chimento and D. Pavón, Phys. Rev. D **73**, 063511 (2006); M.S. Berger and H. Shojaei, Phys. Rev. D **74**, 043530 (2007).
 - [37] G.R. Farrar and P.J.E. Peebles, Astrophys. J. **604**, 1 (2004).
 - [38] J.D. Bekenstein, Phys. Rev. D **49**, 1912 (1994).

- [39] D. Bak and S.-J. Rey, *Class. Quantum Grav.* **17**, L83 (2000); E.E. Flanagan, D. Marolf and R.M. Wald, *Phys. Rev. D* **62**, 084035 (2000).
- [40] B. Guberina, R. Horvat, and H. Nikolic, *JCAP*01(2007)012.
- [41] R. Bean, S.H. Hansen, and A. Melchiorri, *Phys. Rev. D* **64**, 103508 (2001).
- [42] A.R. Liddle and D.H. Lyth, *Cosmological Inflation and Large-Scale Structure* (Cambridge University Press, Cambridge, 2000).
- [43] J.M. Bardeen, P.J. Steinhardt, and M.S. Turner, *Phys. Rev. D* **28**, 679 (1983).
- [44] D.H. Lyth, *Phys. Rev. D* **31**, 1792 (1985).
- [45] J. García-Bellido and D. Wands, *Phys. Rev. D* **53**, 5437 (1996).
- [46] D. Wands, K.A. Malik, D.H. Lyth, and A.R. Liddle, *Phys. Rev. D* **62**, 043527 (2000).
- [47] M.C. Bento, O. Bertolami and A.A. Sen, *Phys. Rev. D* **66**, 043507 (2002).
- [48] W. Zimdahl, arXiv:0705.2131(gr-qc).
- [49] R. Colistete Jr., J.C. Fabris, J. Tossa and W. Zimdahl, arXiv:0706.4086(astro-ph).
- [50] H. Sandvik, M. Tegmark, M. Zaldarriaga and I. Waga, *Phys. Rev. D* **69**, 123524(2004).
- [51] A.B. Balakin, D. Pavón, D.J. Schwarz, and W. Zimdahl, *NJP* **5**, 85.1 (2003).
- [52] R.R.R. Reis, I. Waga, M.O. Calvão, and S.E. Joràs, *Phys. Rev. D* **68**, 061302(2004).
- [53] W. Zimdahl and J.C. Fabris, *Class. Quant. Grav.* **22**, 4311(2005).
- [54] W. Zimdahl, *Int. J. Mod. Phys. D* **14**, 2319 (2005).
- [55] F.C. Carvalho, J.S. Alcaniz, J.A.S. Lima, and R. Silva, *Phys. Rev. Lett.* **97**, 081301 (2006).